

INDORAMIN

What does it do?

Indoramin is used to treat trouble peeing in men caused by an enlarged prostate.

How should you take it?

Take *Indoramin* regularly as directed with a glass of water. Swallow the tablet whole. You can take *Indoramin* with or without food but take it the same way each time.

What if you forget a dose?

Take the missed dose as soon as possible. If it is close to the time for your next dose, skip the missed dose and carry on as normal. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *Indoramin* including:

- Alcohol as this can interfere with the way *indoramin* is absorbed by your body. Drinking alcohol will also increase feelings of sleepiness and dizziness which you may experience with *indoramin*

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Drowsiness, sleepiness	Do not drive or use tools or machines. Do not drink alcohol
Headache, dizziness	Tell your doctor
Changes in sexual function Tiredness or weakness	

Blurred vision Running or blocked nose Stomach upset	Tell your doctor if troublesome
Lightheaded or dizziness after standing up	Stand up slowly. If it is continues, or severe, tell your doctor

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have problems with dizziness, liver or heart problems, or prostate cancer.
- Indoramin can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional.
- If you are having eye surgery (e.g. cataract operation), tell your doctor that you are taking indoramin