

Tamsulosin

What does it do?

Tamsulosin is used to treat trouble peeing in men caused by an enlarged prostate.

How to take it?

Take *tamsulosin* regularly as directed with a glass of water. Swallow the capsules whole. You can take *tamsulosin* with or without food, but take it the same way each time.

What if you forget a dose?

Take the missed dose as soon as possible. If it is close to the time for your next dose, skip the missed dose and carry on as normal. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *tamsulosin* including:

- sildenafil (e.g. Viagra®) or similar medicines

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Chest pain Prolonged erection (longer than four hours)	Tell your doctor immediately
Headache, dizziness	Tell your doctor
Changes in sexual function Tiredness or weakness Blurred vision Runny or blocked nose	Tell your doctor if troublesome

Stomach upset	
Lightheaded or dizzy after standing up	Stand up slowly. If it continues, or is severe, tell your doctor

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have problems with dizziness, liver or heart problems, or prostate cancer.
- *Tamsulosin* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional.
- If you are having eye surgery (e.g. cataract operation), tell your doctor that you are taking *tamsulosin*.